

MTH 4436 HW Set 2.1

FALL 2016

Pat Rossi

Name _____

Set 2.1

1.c. The sum, of any two consecutive triangular numbers is a perfect square.

Note that $t_{n+1} = t_n + (n + 1)$. (Geometrically, the $n + 1^{st}$ triangle is formed by taking the n^{th} triangle and adding a row containing $n + 1$ dots.)

Combine this with the result from part a) that the n^{th} triangular number, $t_n = \frac{n(n+1)}{2}$, and we have:

$$t_n + t_{n+1} = t_n + [t_n + (n + 1)] = 2t_n + n + 1 = 2 \frac{n(n+1)}{2} + n + 1 =$$

$$n(n+1) + n + 1 = (n+1)(n+1) = (n+1)^2$$

$$\text{i.e., } t_n + t_{n+1} = (n+1)^2$$

Hence, the sum of any two consecutive triangular numbers is a perfect square.

7. Show that the difference between the squares of two consecutive triangular numbers is always a cube.

Proof:

Recall: $t_n = \frac{n(n+1)}{2}$ is the n^{th} triangular number.

Consequently: $t_{n+1} = \frac{(n+1)(n+2)}{2}$

Consider: $t_{n+1}^2 - t_n^2 = \left(\frac{(n+1)(n+2)}{2}\right)^2 - \left(\frac{n(n+1)}{2}\right)^2$

$$= \underbrace{\left(\frac{(n+1)(n+2)}{2} + \frac{n(n+1)}{2}\right) \left(\frac{(n+1)(n+2)}{2} - \frac{n(n+1)}{2}\right)}_{\text{Difference of Perfect Squares}}$$

$$= \left(\frac{(n+1)[(n+2)+n]}{2}\right) \left(\frac{(n+1)[(n+2)-n]}{2}\right) = \left(\frac{(n+1)(2n+2)}{2}\right) \left(\frac{(n+1)(2)}{2}\right)$$

$$= \left(\frac{(n+1)(n+1)(2)}{2}\right) \left(\frac{(n+1)(2)}{2}\right) = (n+1)(n+1)(n+1) = (n+1)^3$$

$$\text{i.e., } t_{n+1}^2 - t_n^2 = s_{n+1}^3$$